



# Thermo T7 Transcription Kit

(bacteriophage T7 of *Escherichia coli*)

Cat. No.	size
E0908-01	25 x 25 µl reactions
E0908-02	50 x 25 µl reactions

### Unit Definition:

One unit is the amount of enzyme required to incorporate 1 nmol of labeled UTP into acid-soluble material in 1 hr at 37°C.

### Storage Conditions:

Store at -20°C.

### Quality Control:

All preparations are assayed for contaminating exonuclease, endonuclease, for nonspecific RNase and single- and double-stranded DNase activities. Typical preparations are greater than 90% pure, as judged by SDS polyacrylamide gel electrophoresis.

### References:

1. Chamberlin, M. and Ring, J. (1973) *J. Biol. Chem.* 248, 2235-2244.
2. Tabor, S and Richardson, C.C. (1985) *Proc. Natl. Acad. Sci. U.S.A.* 82, 1074-1078.
3. Sambrook, J., Fritsch, E. F. and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, second edition, pp. 10.27-10.37, Cold Spring Harbour Laboratory, Cold Spring Harbour.

**Thermo T7 Transcription Kit with modified T7 RNA Polymerase for higher efficiency up to 50°C. Extremely useful for radioactive and nonradioactive labeling as well as for RNA synthesis in preparative scale.**

### Description:

- DNA-dependent RNA polymerase with stringent specificity for T7 phage promoters sequence (1) and increased thermostability. Activity range: 37-51°C.
- Ultrapure recombinant enzyme.
- Efficiently synthesizes *in vitro* transcripts from almost any DNA target located downstream from a T7 promoter (2).
- Suitable for preparing labeled single-stranded RNA probes of high specific activity (3).
- Transcripts can be used as hybridization probes, templates for *in vitro* translation, substrates in RNA processing systems, as well as for exon and intron mapping of genomic DNA.

### Storage Buffer:

20 mM potassium phosphate (pH 7.7), 150 mM NaCl, 1 mM EDTA, 1 mM dithiothreitol and 50 % (v/v) glycerol.

### Kit components:

COMPONENT:	E0908-01	E0908-02
5 x T7 Reaction Buffer	150 µl	300 µl
NTPs mix, 25 mM each	50 µl	100 µl
Thermo T7 RNA Polymerase	12.5 µl	25 µl
RNase-free Water	1 ml	1 ml

This product is developed, designed and sold exclusively for research purposes and *in vitro* use only.

EURx Ltd. 80-297 Gdańsk Poland ul. Przyrodników 3, NIP 957-07-05-191, KRS 0000202039  
www.eurx.com.pl, orders@eurx.com.pl, tel. +48 58 524 06 97, fax +48 58 341 74 23

### **In vitro Thermo T7 transcription (25 µl):**

Reaction assembly should be performed at room temperature (not on ice). This prevents any precipitation of template DNA due to spermidine contained in the 5 x T7 Reaction Buffer.

#### 1. Mix:

<b>Component:</b>	<b>Amount:</b>
5 x T7 Reaction Buffer	5 µl
NTPs mix, 25 mM each	2 µl
DNA template*	1-2 µg
Thermo T7 RNA Polymerase**	0.5 µl
RNase-free Water	to 25 µl

2. Incubate up to 2 hours at 37-51°C (the best performance is observed at 46°C), then check transcription on appropriate denaturing polyacrylamide gel.

### **NOTES:**

\*High purity of the template is very important for the yield of reaction. If run off transcription is applied be sure there is no RNase A contamination that could be due to plasmid preparation. We recommend using EURx RNase-free Plasmid DNA Purification Kit (Cat. No. E3500), which works excellent for preparing RNase-free plasmid DNA. In case, T7 template DNA is a PCR fragment, remove primers (recommended: purification from agarose gels using e.g. EURx Agarose-Out DNA Purification Kit, Cat. No. E3540) and confirm DNA homogeneity on an agarose gel.

\*\*0.2 µl of Thermo T7 RNA Polymerase is most efficient for labeling, more enzyme is recommended for preparative scale.