

DNA Polymerase I

(*Escherichia coli*)

Cat. No.	size
E1080-01	500 units
E1080-02	2500 units

Unit Definition: One unit is defined as the amount of enzyme required to incorporate 10 nmoles of total deoxyribonucleotide into acid-insoluble material in 30 minutes at 37°C with DNase I activated DNA as the template primer.

Storage Conditions: Store at -20°C.

References:

1. Lehman, I.R. (1981) *Enzymes* 14, 15-37.
2. Rigby, P.W.J., Diekmann, M., Rhodes, C. and Berg, P. (1977) *J. Mol. Biol.* 113, 237-251.
3. Hartman, C.P. and Robussay, D. (1981) *Gene Amplification and Analysis* (Chirikjian, J.G. and Papas, T.S., eds.) 2, 17-39, Elsevier/North Holland, New York.

DNA Polymerase I is a mezophilic, DNA-dependent DNA polymerase with inherent 3'→5' and 5'→3' exonuclease activity.

Description:

- Exhibits the 5'→3' polymerase activity.
- Exhibits the 5'→3' exonuclease activity, active only on duplex DNA.
- Contains the 3'→5' exonuclease, primarily active on single-stranded DNA (1).
- Ultrapure recombinant enzyme.
- Used to prepare radioactive probes by nick translation (2) and random priming (3).
- Useful for end-labeling of DNA molecules with 3' and 5' protruding tails or blunt-ended.

Storage Buffer:

50 mM potassium phosphate (pH 7.0), 0.25 mM dithiothreitol and 50% (v/v) glycerol.

10 x Reaction Buffer:

500 mM Tris-HCl (pH 7.4 at 25°C), 100 mM MgSO₄, 10 mM dithiothreitol.

Assay Conditions:

67 mM potassium phosphate (pH 7.4), 6.7 mM MgCl₂, 1 mM dithiothreitol, 0.033 mM each dCTP, dGTP, dTTP and [α -³²P]dATP, 4.5 μ g activated DNA. Incubation is at 37°C for 30 min in a reaction volume of 100 μ l.

Quality Control:

All preparations are assayed for contaminating endonuclease, 3'-exonuclease, and nonspecific single- and double-stranded DNase activities. Typical preparations are greater than 95% pure, as judged by SDS polyacrylamide gel electrophoresis.