

TaqII

5'-GACCGA(N)₁₁-3'
3'-CTGGCT(N)₉-5'

Cat. No.	size
E2411-01	100 units
E2411-02	500 units

Reaction Temperature: 65°C

Inactivation Temperature (20 min): –

Prototype: TaqII

Source: *Thermus aquaticus*

Package Contents:

- TaqII
- 10 x ONE Buffer
- Dilution Buffer # TaqII

Added for enzymes exceeding 10 U/μl in concentration. High protein concentration warrants optimal stability during prolonged storage. Use dilution buffer to prepare short term working stocks (5-10 U/μl, non-freezing at -20°C).

Storage Conditions: Store at -20°C.

Double Digestion – Buffer Compatibility:

ONE Buffer is compatible with most EURx restriction enzymes.

Restriction Enzyme Buffer Compatibility:

Both, enzyme and buffers are fully compatible to restrictases and buffer systems from other manufacturers and can be used along in double digestions. To obtain best results, consult the corresponding manuals of all involved products.

DNA Methylation:

No Inhibition: dam, dcm, EcoKI

Potential Inhibition: CpG

Standard Reaction Protocol (for 50 μl volume):

Mix the following reaction components:

1-2 μg pure DNA or 10 μl PCR product (≈0.1-2 μg DNA)

5 μl 10 x ONE Buffer

1-2 U TaqII (use 1 U per μg DNA, < 10% React. Volume!)

Tips: Add enzyme as last component. Mix components well before adding enzyme. After enzyme addition, mix gently by pipetting. Do not vortex. High (excess) amounts of enzyme can greatly speed up the reaction.

add sterile H₂O to 50 μl final volume

Incubate for 3 h at 65°C

Stop reaction by alternatively

- Addition of 2.1 μl EDTA pH 8.0 [0.5 M], final 20 mM *or*
- Heat Inactivation
(not applicable for this enzyme) *or*
- Spin Column DNA Purification
(e.g. EURx PCR/DNA Clean-Up Kit, Cat.No. E3520) *or*
- Gel Electrophoresis and Single Band Excision
(e.g. EURx Agarose-Out DNA Kit, Cat.No. E3540) *or*
- Phenol-Chloroform Extraction or Ethanol Precipitation.

Note 1: It is required to purify DNA before digestion. We recommend PCR / DNA Clean-Up Purification Kit or Agarose-Out DNA Purification Kit.

Note 2: It is not recommended to use more than 1 unit of enzyme per 1 μg of DNA.

Note 3: Over 1 hr digestion is highly recommended. Best results are obtained with 3 hr digestion.

Unit Definition:

One unit is the amount of enzyme required to completely digest unmethylated 1 μg of pBR322 DNA to obtain stable digestion pattern in 1 hr. Total reaction volume is 50 μl. Enzyme activity was determined in the recommended reaction buffer.

Reaction Buffer:

1 x ONE Buffer

Storage Buffer:

20 mM Tris-HCl (pH 7.5 at 22°C), 0.1 mM EDTA, 200 mM NaCl, 1 mM dithiothreitol, 200 μg/ml bovine serum albumin, 0.02% Tergitol™ TMN, 0.02% Tween™20, 50% (v/v) glycerol.

Quality Control:

All preparations are assayed for contaminating endonuclease, 3'-exonuclease, 5'-exonuclease/5'-phosphatase, as well as nonspecific single- and double-stranded DNase activities.